

### Things you can do on your own

#### Keep track of your diarrhea:

• Write down how many bowel movements you have each day

### Stay hydrated and change your diet to lessen your symptoms:

- Drink at least 6-8 cups of clear liquids, such as water, daily to prevent dehydration
- Avoid caffeine, alcohol, milk, and artificial sweeteners
- Avoid greasy, fatty, and fried foods
- Eat small, frequent meals throughout the day instead of 3 big meals
- Eat foods that help diarrhea like bananas, melon, rice, applesauce, and/or toast
- Drink sports drinks such as Gatorade to replenish your potassium

### Keep the skin around your anus healthy:

- If your skin is getting irritated after bowel movements, gently wipe yourself with baby wipes and then apply barrier creams, such as zinc oxide, Vaseline<sup>®</sup>, or diaper ointments
- Add 1-2 tablespoons of baking soda to a warm tub of water and soak your bottom

See the Other Resources page for more information

### With over-the-counter medications

- IF YOU ARE TAKING an IMMUNOTHERAPY DRUG (i.e., Keytruda/pembrolizumab, Opdivo/nivolumab, Libtayo/cemiplimab, Tecentriq/atezolizumab, Bavencio/avelumab, Imfinzi/durvalumab or Yervoy/ipilimumab), DO NOT TAKE OVER THE COUNTER MEDICATIONS for diarrhea.
- Several treatments for diarrhea, such as Imodium, are available over the counter at your local pharmacy. Please use these medicines as directed.

**REMINDER** - Always keep your care team updated about over-the-counter medicines you are taking. If your care team tells you they are not safe for you, do not take them.

This content was created for the SIMPRO Study and is provided as general health information and is not intended to: invite or establish a healthcare providerpatient relationship, constitute furnishing professional services, constitute, or substitute for, the advice or judgment of a medical professional; or serve as a basis for medical treatment. Updated: September 2022.

© 2022 SIMPRO Study Team (Dana-Farber Cancer Institute Inc., Baptist Memorial Health Care Corporation, Dartmouth-Hitchcock Medical Center, Lifespan Cancer Institute, West Virginia University Cancer Institute, Maine Medical Center). All Rights Reserved.

# With the help of your care team

If over-the-counter medications are not helping, your care team may give you a prescription medicine, such as:

- Lomotil (or atropine-diphenoxylate)
- If you have trouble with pills, tincture of opium or "DTO" may be an option

# When to call your care team for help

- If you have a fever ≥100.5°F (38°C) along with the diarrhea
- If your stools are watery
- If your diarrhea gets worse, not better, after following these tips
- If you are having signs of dehydration. Signs of dehydration include:
  - o Feeling weak, dizzy, or lightheaded
  - Having dark yellow or very little urine
  - Having a dry mouth/being thirsty
- If you are waking up in the middle of the night with diarrhea
- If you have an accident, or lose control of your bowels
- If there is blood in your stool
- If your stools are black in color, as this can sometimes be a sign of internal bleeding

# Additional information on Diarrhea

- <u>Click here</u> for more information on managing diarrhea
- <u>Click here</u> for how cancer can affect diarrhea

© 2022 SIMPRO Study Team (Dana-Farber Cancer Institute Inc., Baptist Memorial Health Care Corporation, Dartmouth-Hitchcock Medical Center, Lifespan Cancer Institute, West Virginia University Cancer Institute, Maine Medical Center). All Rights Reserved.