



Tips for Managing Nausea and Vomiting

Things you can do on your own



Understand what might be making you feel sick:

- Keep a diary tracking when your nausea is worst in your chemotherapy cycle
- Record what triggers your nausea
- Make sure you understand what medicines to take, and when to take them

Eat and drink well:

- Eat small, frequent meals, and supplement with meal replacement drinks, such as Boost or Ensure
- Drink 6-8 cups of fluids, such as water, per day
- Avoid greasy or fatty foods, like fried foods
- Eat bland foods
- Try sucking on hard candies made with lemon or ginger
- Stay away from strong smells

Get comfortable:

- Wear loose-fitting clothes
- Do mind-body exercises, or try deep breathing exercises
- Try acupuncture

See the Other Resources page for more information

With the help of your care team



- Your care team might give you a prescription medication, and you might have to try a few different nausea medicines before you find the combination that works best for you
- Your care team might recommend that you take your anti-nausea medication:
 - On a regular schedule to prevent nausea, even if you aren't feeling nauseated at the moment
 - Only when you feel nauseated (usually written "as needed" on the prescription bottle)
 - Some medications on a regular schedule, and others as needed
- You may find that you need to take medications on particular days of your chemotherapy cycle
- If you are not able to keep your anti-nausea medication down, then you can receive them in a different way, such as by IV, suppository, or dissolving tablet

This content was created for the SIMPRO Study and is provided as general health information and is not intended to: invite or establish a healthcare provider-patient relationship, constitute furnishing professional services, constitute, or substitute for, the advice or judgment of a medical professional; or serve as a basis for medical treatment. Updated: September 2022.

© 2022 SIMPRO Study Team (Dana-Farber Cancer Institute Inc., Baptist Memorial Health Care Corporation, Dartmouth-Hitchcock Medical Center, Lifespan Cancer Institute, West Virginia University Cancer Institute, Maine Medical Center). All Rights Reserved.



When to call your care team for help



- If you are taking anti-nausea medication as prescribed and it is not working, or it stops working
- If there is blood, or something that looks like coffee grounds in your vomit
- If you cannot keep anything (food, water, or medicines) down because of your vomiting
- If you go a day without drinking any fluids
- If you have not passed any gas for a day or more
- If you have not had a bowel movement for 2 or more days
- If you have severe nausea or vomiting and new abdominal (belly area) pain
- If you are dehydrated, or don't have enough fluid in your body. Signs of dehydration include:
 - Feeling weak, dizzy, or lightheaded
 - Having dark yellow or very little pee
 - Having a dry mouth/being thirsty

****When you call it helps to have a diary listing what medicines you have been taking and notes about how much you have been able to drink to stay hydrated****

Additional information on Nausea and Vomiting



- [Click here](#) for a video about cancer related nausea and vomiting